

2000 Annual Report on Work-Related Asthma in Michigan



2000 Annual Report on Work-Related Asthma in Michigan

A Joint Report
of the
Michigan State University
Department of Medicine
117 West Fee Hall
East Lansing, Michigan 48824-1315
(517) 353-1846
Kenneth D. Rosenman, MD
Professor of Medicine
Mary Jo Reilly, MS
Epidemiologist

and

the Michigan Department
of Consumer & Industry Services
P. O. Box 30649
Lansing, Michigan 48909-8149
(517) 332-1608
Douglas J. Kalinowski, MS, CIH
Deputy Director
Bureau of Safety and Regulation

June 8, 2001

Summary:

This is the tenth annual report on work-related asthma (WRA) in Michigan. An average of 145 new people each year are reported to the Michigan Department of Consumer and Industry Services (MDCIS) with asthma caused by work. The reports received by the Michigan Department of Consumer and Industry Services represent 18.5%-64.9% of the number of estimated people in Michigan who develop work-related asthma each year. Inspections at the workplaces of these individuals reveal a large number of fellow workers with asthma or respiratory symptoms compatible with asthma. The work exposures may cause new onset asthma from exposure to an allergen or an irritant that precipitates inflammatory changes, or work exposures may exacerbate pre-existing asthma from exposure to an irritant. Almost ninety percent of the reports received involve the new onset of asthma.

Known allergens such as isocyanates and metal working fluids are the most common cause of work-related asthma in Michigan. Despite the association of a patient's asthma with work, many times a known allergen is not identified. Practitioners need to maintain a high level of clinical suspicion for new causes of work-related asthma. There are approximately 350 documented agents associated with work-related asthma. The most comprehensive listing of known causes of work-related asthma is at an internet website:
www.remcomp.com/asmanet/asmapro/asmawork.htm#star.

Work-related asthma is affecting men and women, generally in the 30 - 50 year old age range. The average annual incidence rate of work-related asthma among African Americans was 2.1 times greater than among whites. Exposures are occurring in a wide range of workplaces. When an inspection is conducted at the workplace, significant numbers of symptomatic individuals have been identified. However, air monitoring at these same facilities typically reveals that the exposures to the suspected allergen or irritant are within existing workplace standards. Often a specific Michigan Occupational Safety and Health Act (MIOSHA) workplace standard for the suspected allergen or irritant does not exist.

In 1998, the National Institute for Occupational Safety and Health (NIOSH) recommended reducing metal working fluid aerosols to 0.4 mg/m³ (thoracic particulate mass) as a time-weighted average (TWA) for up to 10 hours/day for a 40 hour work week. In addition to lowering the recommended exposure limit (REL), NIOSH recommends the development of a comprehensive health and safety program at companies that use metal working fluids. NIOSH recommends that the program include: 1) safety and health training; 2) worksite analysis; 3) hazard prevention and control, and; 4) medical monitoring of exposed workers (1). The Federal Occupational Safety and Health Administration (OSHA) is considering but has not yet proposed adopting these recommendations as an enforceable standard.

This coming year, we will be working on two new initiatives concerning work-related asthma. Three states in addition to Michigan have developed surveillance systems for work-related asthma.

An in-depth analysis of information related to cleaning agent exposures that the four states doing surveillance of WRA have collected is underway. By combining the data on cleaning agent exposures from the four states doing work-related asthma surveillance, California, Massachusetts, Michigan and New Jersey, we will be able to determine the characteristics and trends of workers who develop asthma from these types of exposures. The second initiative we will be working on this coming year is a Local Emphasis Program (LEP) on workers exposed to welding fume. We will be collecting and analyzing information obtained during MIOSHA enforcement inspections to better understand the types of welding fume exposures and work practices associated with WRA. These two initiatives were undertaken because these exposures account for a significant amount of the work-related asthma cases identified through our surveillance efforts.

Background:

Based on extrapolation from national estimates there are approximately 230,000 adults with asthma in Michigan (2). It is estimated that work place exposures are responsible for asthma in 2 - 33% of adults (3-12). One Michigan study estimated that workplace exposures are possible contributing causes to the development of asthma in 3 -26% of adults hospitalized with asthma in Michigan (6). A recent review of the literature estimated that 2.8% - 21% of asthma in adults is related to work (12).

In 1988, the Michigan Department of Consumer and Industry Services (formerly the Michigan Department of Public Health) instituted a surveillance program for work-related asthma with financial assistance from the National Institute for Occupational Safety and Health (NIOSH). The goal of the surveillance program is to prevent work-related asthma through the reporting of index patients. The reporting of the index patient is regarded as a sentinel health event that may lead to the identification of other employees from the same facility who are at risk of developing asthma or who have developed similar breathing problems. There are three sources used to identify persons with work-related asthma: reports from physicians; reports from hospitals (since 1989); and claims filed since 1988 with the Bureau of Workers' Compensation. Both physicians in private practice and those working for industry send reports to the Michigan Department of Consumer and Industry Services (MDCIS). Reports from hospitals are requested once each year. Hospital discharge summaries for individuals with a primary or secondary diagnosis of a respiratory condition due to chemical fumes and vapors (ICD-9 506.0-.9) as well as individuals with a primary or secondary diagnosis of asthma (ICD-9 493) where the principal payer is listed as workers' compensation are obtained from the Michigan Health and Hospital Association's (MHA) Michigan inpatient database to verify the completeness of reporting by the hospitals, and the work-relatedness of the condition is determined.

A person is considered to have work-related asthma from sensitization to a workplace exposure if: A) they have a physician diagnosis of asthma, and B) onset of respiratory symptoms associated

with a particular job that then improve or are relieved when the patient is not working, and C) they work with a known occupational allergen, or have evidence of an association between work exposures and a decrease in pulmonary function testing.

Additional criteria are used to record three other categories of asthma associated with work. If only criteria (A) and (B) above are met the person is considered to have possible work-related asthma. An industrial hygiene investigation at the patient's work site is performed to determine the allergen and to document its associated exposure levels. If a person had physician-diagnosed asthma before beginning work and their asthma became worse at a particular job the person is considered to have work-aggravated asthma. Occupational asthma from exposure to an allergen at work typically develops after a variable period of symptomless exposure to the sensitizing agent. However, if a person develops asthma for the first time immediately after an acute exposure to an irritating chemical at work the patient is considered to have Reactive Airways Dysfunction Syndrome (RADS) (13).

After the patient has been interviewed and the work-relatedness of the condition evaluated, an industrial hygiene investigation may be conducted at the patient's workplace. At this follow-up investigation, co-workers are interviewed to determine if other individuals are experiencing similar breathing problems from exposure to the suspected allergen. An industrial hygienist conducts air monitoring for any suspected allergens and reviews the company's health and safety program. After the investigation is completed, a report of air sampling results and any recommendations are sent to the company and union (or designated labor representative, if the company does not have a union).

In 1998, the surveillance program also began sending letters about potential problems with exposures to some of the companies where the index cases were exposed to an allergen, in cases where a formal MIOSHA inspection was not planned. These letters, in lieu of inspections, are sent to the company health and safety director, and ask the company to evaluate exposures to whatever suspected allergens were identified through the telephone interview with the index case.

Results:

Reports

Table 1 shows that 1,638 people were confirmed with work-related asthma between 1988 - 2000. Figure 1 presents the same data in a bar graph of the number of cases identified each year, and the types of work-related asthma that were confirmed. The reports are divided into four categories: occupational asthma, possible occupational asthma, aggravated asthma and RADS. One hundred forty-one additional patients have been confirmed since last year's report. There were 31 confirmed reports in 1988, 63 in 1989, 144 in 1990, 115 in 1991, 150 in 1992, 176 in

1993, 152 in 1994, 127 in 1995, 155 in 1996, 162 in 1997, 147 in 1998, 119 in 1999, and 97 in 2000. Figure 2 shows the overlap of the 1,541 patients by reporting sources for 1988-1999.

The data is incomplete for 1988 since the surveillance system was initiated in that year. To date we have not yet received complete hospital reporting for the years 1999 and 2000. Patient interviews are still needed for 6 reports of patients from hospitals and 4 reports of patients from physicians in 1999; and, for 14 reports of patients from physicians in 2000.

Gender

Eight hundred forty (51.3 %) of the persons with work-related asthma are women and 798 (48.7 %) are men.

Race

One thousand two hundred fifty-two (76.8 %) of the persons with work-related asthma are white, 305 (18.7 %) are African American, 41 (2.5 %) are Hispanic, 15 (0.9 %) are Alaskan or American Indian, 6 (0.4 %) are Asian and, 12 (0.7 %) were listed as "other." For seven individuals, race was unknown.

The average number of incident cases of African Americans with work-related asthma each year for 1992-1998 was 29.7. In 1998 there were 539,621 African Americans in the Michigan labor force (14). The annual incidence rate for work-related asthma in African Americans, therefore, was 5.5/100,000 workers. The respective data for whites was 115.3 new cases per year and 4,368,720 whites in the Michigan labor force. The annual incidence rate of work-related asthma in whites, therefore, was 2.6/100,000 workers. The African American rate of work-related asthma was 2.1 times greater than the rate for whites.

Age

The dates of birth range from 1905 - 1978. The average year of birth is 1952.

Location in State

Figure 3 shows the county in which the patient worked where they developed work-related asthma. The main locations are: Wayne (424 cases, 26.3 %), Oakland (219 cases, 13.6 %), and Macomb (160 cases, 9.9 %). Table 2 and Figure 4 show the annual average incidence rates of work-related asthma among the general working population in each county. Based on the annual average incidence of reports of confirmed cases per 100,000 adult workers, Luce (19.5 per 100,000), Cheboygan (11.9 per 100,000) and Clare (11.2 per 100,000) have the highest rates. It should be noted that, even though Luce county had the highest incidence rate of work-related asthma, the rate is based on only four cases (see Table 2). Table 3 shows the annual incidence

rates for the larger metropolitan areas and the whole state for the years 1990 through 1998 separately.

Type of Industry

Table 4 shows the types of Michigan industries where the exposures to the occupational allergens occurred from 1988 to 2000. Figure 5 shows the distribution of major industry types for all cases identified from 1988-2000. The predominant industries for the total number of cases identified between 1988 and 2000 were in the manufacturing sector: automobile (43.3%), fabricated metal products (4.4%), rubber and miscellaneous plastic products (4.0%), industrial and commercial machinery and computer equipment (3.8%), and foundries (3.4%). Workers in the health field also accounted for a high percentage of the total number of patients (9.0%).

The incidence rate of work-related asthma by industry type ranges from 0.3 cases per 100,000 in several industries (restaurants, general merchandise stores and farming) to a high of 20.9 cases per 100,000 in the manufacture of transportation equipment. The industries with the highest annual average incidence rates besides the manufacture of transportation equipment included: foundries with 13.3 cases per 100,000 workers; education with 11.8 cases per 100,000 workers; the manufacture of other nondurables with 11.4 cases per 100,000 workers; and the manufacture of rubber products with 10.5 cases per 100,000 workers.

Table 5 shows the annual incidence rates for the 1990 through 1998 work-related cases within those industries which had 20 or more reports.

Overall, by broad industrial classification, the average annual incidence rates were: 11.0 cases per 100,000 workers in the manufacturing industry; 2.7 cases per 100,000 workers in the construction and mining industry; and 0.8 cases per 100,000 workers in the service producing industry.

Table 6 shows the predominant exposures causing work-related asthma in Michigan. The most frequent exposures were to isocyanates (18.0%), metal working fluids (12.1%), vehicle exhaust (5.4%), welding fumes (5.0%), and solvents (3.0%). The agent has not yet been identified for 207 patients (12.6%). The exposures to unknown agents occurred 107 times in the manufacturing sector and 100 times in an office setting.

Medical Results

Table 7 shows patients' cigarette smoking status. Almost 20% of patients were smoking when their work-related asthma developed. This is a lower percentage than the state average and markedly lower than that found in a blue collar working population.

Forty-two percent of the work-related asthma patients had a family history of allergies (Table 8).

Forty-five percent of the asthma patients had a personal history of allergies or asthma (Table 9). Two hundred ninety-three (41.3%) of the 710 patients with a personal history of allergies or asthma previously had asthma.

One thousand three hundred twenty-six of the patients identified with work-related asthma had persistence of their asthma symptoms (Table 10). This was true for 442 of 464 (95.3%) of those still exposed as well as 884 of 1057 (83.6%) no longer exposed to the substance causing their asthma. Among those no longer exposed, 50.0% stated their symptoms were less severe compared to 33.2% among those still exposed who reported their symptoms were less severe. Similarly, 84.5% of those still exposed were continuing to take asthma medications while 76.6% of those no longer exposed were still taking asthma medications. Among those no longer exposed, 29.9% stated they were taking fewer medications while only 21.1% of those still exposed were taking fewer medications (Table 10).

Seven hundred nine of 1,494 (47.5%) patients with known workers' compensation status had applied for workers' compensation. Cases were pending for 361 (50.9%) of those who applied, while 245 (34.6%) had received awards and 103 (14.5%) had been denied.

Although 1,638 individuals were confirmed with work-related asthma, we could find objective testing for hyper-reactivity by methacholine challenge for 25% of cases, by specific bronchoprovocation for 1% of cases, and by pre- and post- bronchoprovocation for 53% of cases. In addition, we found only 3% of cases had peak flow monitoring and only 2% of cases had pre- and post-work shift testing.

Industrial Hygiene

The 1,638 people with work-related asthma worked at 1,089 different facilities. Another 62 facilities were inspected more than once. Inspections were performed at 513 (44.6%) of these 1,151 facilities. Thirty-eight inspections were completed since last year's report. Inspections are scheduled at 3 (0.3%) facilities (Table 11). In addition, because of inadequate resources to conduct inspections at all the identified facilities, 62 facilities received letters notifying them that a disease report had been received and asked them to investigate potential exposures causing the respiratory problem, including indoor air problems.

Air sampling for allergens was conducted during 362 of the inspections. Forty-nine of the 362 (13.5%) facilities were above the National Institute for Occupational Safety and Health recommended exposure limit (REL). Twenty-four of the 362 (6.6%) were above the enforceable Michigan Occupational Safety and Health Act (MIOSHA) permissible exposure limit (PEL) (Table 12).

Interviews of fellow workers were performed at 389 of the 513 inspections. Workers had daily or weekly breathing symptoms or onset of new asthma since beginning to work at that company

in 278 of the 389 (71.5%) companies. The average percentage of workers with symptoms in these 278 companies was 21.0%, ranging from 2% to 100%. Interviews conducted in 111 companies found no co-workers with symptoms. One thousand two hundred and fifty-seven of the 7,337 (17.1%) fellow workers interviewed had symptoms consistent with work-related asthma (new onset asthma or bothered at work by daily or weekly shortness of breath, wheezing or chest tightness) (Table 13).

Four hundred thirty workers from 90 companies were listed by employers on the Occupational Safety and Health Administration (OSHA) Injury and Illness log as having asthma or asthma-like symptoms. Eight workers had both daily or weekly breathing symptoms and were also listed on the OSHA log. Therefore, a total of 1,679 symptomatic workers were identified during the 513 inspections.

Comparison of Work-Related Asthma in Michigan with Other Geographic Areas

The annual incidence rate of 3.4 cases of work-related asthma per 100,000 workers in the Michigan surveillance system is similar to the rates reported in other surveillance systems. In England the rates reported vary from 2.0 to 4.3 cases per 100,000 with rates as high as 6.3 per 100,000 reported in some regions (15-17). Similarly in Michigan there is regional variation with incidence rates in some counties as high as 19.5 cases per 100,000, although this high rate was based on a small number of cases (Table 2). Reported rates in specific occupations in England have ranged as high as 65.0 cases per 100,000 (17). The system in England is voluntary, and is based only on reports received from pulmonary and occupational physicians (18). This differs from Michigan's surveillance system which is mandatory and covers physicians of all specialties. In Michigan 49% of the reports are from occupational and pulmonary physicians, 7% are from allergists and the remaining 44% are from other specialties. Finland has a national mandatory reporting system that is connected with their workers' compensation system; their reported rate is 15.2 cases per 100,000 (19). Isocyanates are typically the most common cause of work-related asthma reported in other surveillance systems (18,20). However, since farmers were included in the workers' compensation system in Finland, cow dander has become the most commonly reported allergen in that country (21).

A more recent report from a study performed at an HMO in Massachusetts estimated the annual rate of work-related asthma at 71.0 cases per 100,000 (11). Possible reasons for this much higher rate reported in the study from Massachusetts include: 1) less restrictive case criteria in the HMO study; 2) more intense case-finding in the HMO study; 3) a general increase in the incidence of work-related asthma between the time of the more recent HMO study and previous studies; and 4) the fact that the annual estimates from the HMO study were extrapolated from cases identified during the 3 month period of July, August, and September.

Discussion:

In our previous annual reports, we have emphasized the fact that the cases reported in Michigan's surveillance system are likely an under count of the true number of cases of work-related asthma in the state. This continues to be true. Studies suggest that work exposures are important etiologic agents in a significant percentage (20% or greater) of adults with asthma (3-12).

An average of 145 new people each year are reported to the Michigan Department of Consumer and Industry Services (MDCIS) with confirmed work-related asthma. National estimates would project that there are a total of 4,600 - 34,500 adults with work-related asthma in the state (3-12). A Michigan study predicts possibly 6,900 - 59,800 adults whose work contributed to the development of their asthma (6). Using capture-recapture analysis we estimate 228 - 801 adults in Michigan develop work-related asthma each year (12).

As in the previous annual reports on work-related asthma in Michigan, the workers reported are generally young to middle age white men and women, with the greatest number being reported from the Detroit metropolitan area. However, the rate of work-related asthma in African Americans is 2.1 times greater than among whites. The people typically develop their asthma from exposures to agents in the manufacturing sector, particularly automobiles, machinery, metals, chemicals, and rubber and plastics. The predominant causes of work-related asthma remain isocyanates and metal-working fluids.

Asthma symptoms persist despite removal from the precipitating work exposures (Table 10). Studies have shown that the sooner an individual is removed from the exposure causing their asthma after symptoms develop the more likely the individual's symptoms will resolve. On the average among the 1057 individuals no longer exposed, 2.7 years elapse from time of onset of respiratory symptoms at work to date last exposed. We do not have data on how much of this delay is secondary to the individual not seeking medical care and how much is related to the physician not recommending that the individual leave the exposure.

Neither personal habits such as cigarette smoking nor individual susceptibility as measured by personal or family history of allergies are predictive of who will develop work-related asthma. Approximately 50% of the asthma patients have no personal or family history of allergies and 80% are not smoking cigarettes at the time their asthma symptoms develop (Tables 7-9).

Although most facilities where the patient developed asthma were not in violation of exposure standards, there were high percentages of symptomatic fellow workers in facilities using occupational allergens. We identified 1,257 fellow workers with symptoms compatible with work-related asthma. Four hundred thirty individuals were listed on the OSHA log as having work-related asthma. There was only an overlap of eight individuals. Half of the symptomatic individuals indicate they have never seen a doctor for their respiratory symptoms. This indicates the need for more companies to implement medical surveillance programs. The high percentages

of symptomatic individuals are consistent with estimates of the prevalence of work-related asthma in the state. The presence of symptomatic co-workers suggests that some of the occupational health standards may not be sufficiently protective to ensure a safe workplace.

Reevaluations of some of the allowable exposure standards may be needed. These reevaluations might consider comprehensive workplace controls which address not only daily average exposures but exposures during spills and leaks and provision of medical surveillance for potentially exposed workers. Medical monitoring is particularly relevant to reducing the burden of work-related causes of asthma. The longer a person with symptoms remains exposed, the more likely their asthma will become a chronic problem (22). The percentages of individuals reported with work-related asthma for whom this surveillance system could document as having had breathing tests performed in relation to work is less than 10%. This reflects the standard of medical care in the United States where the diagnosis of work-related asthma is made from patient history. More frequent use of objective pulmonary function testing performed in relation to work would allow health care providers to feel more confident about advising their patients to leave their exposure. Cessation of exposure is the most important aspect of treatment of work-related asthma. Patients who are removed from exposure the soonest have the best prognosis (22).

The Michigan Department of Community Health is preparing a planning document for a new effort to reduce Michigan's asthma burden, the Michigan Asthma Strategic Planning Initiative (MASPI). Work-related asthma needs to be integrated into new overall asthma initiatives planned on surveillance and education, both for health care providers and the public.

We are planning new initiatives for this coming year on cleaning agents and welding fumes, two of the most common causes of work-related asthma in Michigan. The cleaning agent initiative will first analyze the data from the four states currently conducting surveillance for work-related asthma: California, Massachusetts, Michigan and New Jersey. After an examination of the characteristics and trends associated with cleaning agent exposures, we plan to develop educational materials that could help prevent cleaning agent exposures that lead to work-related asthma, and distribute this information to stake holders in a position to work toward this goal.

The second planned initiative involves the development and implementation of a Local Emphasis Program (LEP) for welding fumes. This program will work to identify workplaces with welding fume exposures during MIOSHA enforcement inspections. Collection of information during these inspections will allow us to better understand the factors most associated with the development of breathing problems and work-related asthma from welding fume exposures. Both of the initiatives planned for this coming year with further our goal toward reducing the burden of occupational diseases in Michigan.

The report of a patient with known or suspected work-related asthma is a sentinel health event that is critical to effective occupational disease surveillance. Case reporting from physicians offers the opportunity for the most timely workplace interventions, compared to receiving reports from hospitals. With continued support and increasing awareness of work-related asthma by physicians

and other health professionals, we can continue to provide timely intervention in the workplace, offer suggestions for reducing workplace exposures even if they are below current permissible exposure levels, document the need for the development of new standards, identify new occupational allergens and prevent co-workers from developing disease.

References

1. National Institute for Occupational Safety and Health (NIOSH). *Criteria for a Recommended Standard: Occupational Exposure to Metal Working Fluids*. (NIOSH) Publication No. 98-102.
2. *Guidelines for the Diagnosis and Management of Asthma*. NHLBI, National Asthma Education Program Expert Panel Report. *Journal of Allergy and Clinical Immunology* 1991; 88: 425-534.
3. Kobayashi S. *Different Aspects of Occupational Asthma in Japan*. In: Frazier CA, ed 1980 Occupational Asthma. New York: Van Nostrand Reinhold, 1980; 229-244.
4. Blanc P. *Occupational Asthma in a National Disability Survey*. *Chest* 1987; 92:613-617.
5. Salvaggio J ed. *Occupational and Environmental Respiratory Disease in NIAID Task Force Report: Asthma and Other Allergic Disease*. Washington D.C.: U. S. Department of Health Education and Welfare, May 1979 (NIH Publication no. 79-387).
6. Timmer S, and Rosenman KD. *Occurrence of Occupational Asthma*. *Chest* 1993; 104:816-820.
7. Reijula K, Haahtela T, Klaukka T and Rantanen J. *Incidence of Occupational Asthma and Persistent Asthma in Young Adults has Increased in Finland*. *Chest* 1996; 110:58-61.
8. Ng TP, Hong CY, Goh LG, Wang ML, Koh KTC and Ling SL. *Risks of Asthma Associated with Occupations in a Community-Based Case-Control Study*. *American Journal of Industrial Medicine* 1994; 25:709-718.
9. Kogevinas M, Anto JM, Soriano JB, Tobias A and Burney P. *The Risk of Asthma Attributable to Occupational Exposures. A Population-Based Study in Spain*. *American Journal Respiratory and Critical Care Medicine* 1996; 154:137-143.
10. Blanc PD, Cisternas M, Sarith S and Yelin EH. *Asthma, Employment Status, and Disability Among Adults Treated by Pulmonary and Allergy Specialists*. *Chest* 1996; 109:56S-57S.

11. Milton DK, Solomon GM, Rosiello RA, Herrick RA. *Risk and Incidence of Asthma Attributable to Occupational Exposure Among HMO Members*. American Journal of Industrial Medicine 1998; 33:1-10.
12. Henneberger PK, Kreiss K, Rosenman KD, Reilly MJ, Chang YF, Geidenberger CA. *An Evaluation of the Incidence of Work-Related Asthma in the United States*. International Journal of Occupational and Environmental Health 1999; 5:1-8.
13. Brooks SM, Weiss MA, and Bernstein IL. *Reactive Airways Dysfunction Syndrome(RADS)*: Persistent Asthma Syndrome After High Level Irritant Exposures*. Chest 1985; 83:376-384.
14. Michigan Department of Career Development (MDCD), Employment Service Agency. *Affirmative Action Information Report: Michigan 1999*. Detroit, MI. Office of Labor Market Information
15. Gannon PFG, Burge PS. *A Preliminary Report of a Surveillance Scheme of Occupational Asthma in the West Midlands*. British Journal Industrial Medicine. 1991; 48:579-582.
16. Meredith SK, Taylor VM, McDonald JC. *Occupational Respiratory Disease in the United Kingdom 1989: A Report to the British Thoracic Society and the Society of Occupational Medicine by the Project Group*. British Journal Industrial Medicine. 1991; 48:292-298.
17. Meredith S. *Reported Incidence of Occupational Asthma in the United Kingdom, 1989-1990*. Journal Epidemiol Community Health. 1993; 47:459-463.
18. Gannon PFG, Burge PS. *The SHIELD Scheme in the West Midlands Regions, United Kingdom*. British Journal Industrial Medicine. 1993; 50:791-796.
19. Keskinen H. *Epidemiology of Occupational Lung Disease: Asthma and Allergic Alveolitis*. In: Kerr JW, Ganderton MA, eds. *Proceedings of XI International Congress of Allergy and Clinical Immunology*, London, 1982. London: MacMillan; 1983:403-407.
20. Lee HS, Phoon WH, Wang YT, et al. *Occupational Asthma in Singapore. A Review of Cases from 1983 to 1990*. Singapore Medical Journal. 1991; 32:398-402.
21. Reijula K, Patterson R. *Occupational Allergies in Finland in 1981-1991*. Allergy Proc. 1994; 15:163-168.
22. Chan-Yeung M and Malo JL. *Occupational Asthma*. New England Journal of Medicine 1995; 333:107-112.

This report was funded by the National Institute for Occupational Safety and Health under cooperative agreement #U60-CCU502998-14.

**Figure 1. Number of Confirmed Cases of Work-Related Asthma
by Year and Type**






Figure 2. Overlap of Reporting Sources for Confirmed Work-Related Asthma Patients: 1988-1999*



* Diagram represents 1,541 individuals initially reported from 1988-1999. N's represent the total number for that source. Numbers in parentheses represent the overlap of reporting sources. Reporting Source Codes: HDC=Hospital Discharge Data; PR=Physician Referral; DL=Department of Labor; ICFU=Index Case Follow-Up; MSHA=Mine Safety and Health Administration.


Figure 3. Distribution of Confirmed Work-Related Asthma Patients by County of Exposure: 1988-2000



Oakland and **Wayne** counties had the highest number of work-related asthma patients, with 219 and 424 individuals, respectively.


*County of exposure was unknown for 10 patients. Eighteen patients were exposed to an allergen out-of-state.

Figure 4. Average Annual Incidence Rates of Work-Related Asthma Among Michigan Workers by County of Exposure: 1989-1998*



*Rate per 100,000 among Michigan workers. Source: MESC 1993 Annual Average Labor Statistics for Employment by Place of Work. In 1993, there were a total of 3,982,000 Michigan workers.

**Figure 5. Major Industry Type for
Confirmed Work-Related Asthma Patients: 1988-2000***



*Number of patients in parentheses, percent below.

Table 1. Number of Confirmed Cases of Work-Related Asthma by Year and Type

Year	Disease Status*				<u>TOTAL</u>
	<u>OA</u>	<u>POA</u>	<u>AA</u>	<u>RA</u>	
1988	23	7	0	1	31
1989	43	12	3	5	63
1990	87	35	14	8	144
1991	55	30	14	16	115
1992	81	37	14	18	150
1993	75	69	13	19	176
1994	65	59	15	13	152
1995	56	35	19	17	127
1996	60	60	24	11	155
1997	53	74	19	16	162
1998	44	76	18	9	147
1999	44	58	11	6	119
2000	36	42	13	6	97
Total	722	594	177	145	1638

*OA = occupational asthma; POA = possible occupational asthma; AA = aggravated asthma; RA = reactive airway dysfunction syndrome.

**Table 2. Average Annual Incidence Rates of Work-Related Asthma
Among Michigan Workers by County of Exposure: 1989-1998**

<u>County</u>	<u>Number Of Employees*</u>	<u>Avg. Ann. Inc. Rate**</u>	<u>Total # Cases 1989-1998</u>
Alcona and Iosco (0, 2 cases respectively)	9,750	2.1	2
Alpena	11,925	2.5	3
Antrim	4,900	4.1	2
Arenac	4,175	4.8	2
Baraga	2,975	3.4	1
Berrien	67,100	2.5	17
Branch	11,900	6.7	8
Cass	10,000	3.0	3
Charlevoix	9,075	3.3	3
Cheboygan	6,725	11.9	8
Chippewa	13,825	2.2	3
Clare	7,150	11.2	8
Clinton-Eaton-Ingham (4, 9, 42 cases respectively)	213,700	2.6	55
Crawford	4,475	4.5	2
Delta	13,825	1.4	2
Dickinson	12,925	5.4	7
Emmet	13,550	1.5	2
Genesee	164,400	5.8	96
Gladwin	4,475	2.2	1
Grand Traverse-Benzie-Kalkaska-Leelanau (8, 2, 5, 2 cases respectively)	51,525	3.3	17
Gratiot	13,475	2.2	3
Hillsdale	13,925	5.0	7
Huron	11,325	7.1	8
Ionia	14,875	5.4	8
Iron	3,800	2.6	1
Isabella	21,675	1.4	3
Jackson	55,200	4.3	24
Kalamazoo-Calhoun -VanBuren (12,13,3 cases respectively)	195,400	1.4	28
Kent-Ottawa-Muskegon -Allegan (34, 13, 15, 12 cases respectively)	464,800	1.6	74
Luce	2,050	19.5	4
Manistee	6,850	1.5	1
Marquette	28,250	1.8	5
Mason	9,850	1.0	1
Menominee	8,825	1.1	1
Montcalm	17,375	5.2	9
Montmorency	2,200	4.5	1
Newaygo	9,300	3.2	3
Ogemaw	5,575	1.8	1
Osceola	7,050	2.8	2
Otsego	8,850	9.0	8
Sanilac	11,400	4.4	5
Shiawassee	18,875	2.1	4
St. Joseph	22,000	1.8	4
Tuscola	13,250	6.8	9
Washtenaw-Lenawee-Livingston (86, 12, 20 cases respectively)	244,700	4.8	118
Wexford-Missaukee (3, 0 cases respectively)	15,300	2.0	3
Saginaw-Bay-Midland (35, 10, 7 cases respectively)	164,600	3.2	52
Detroit MSA***	1,877,000	3.9	738
Out of State	--	--	17
Unknown	--	--	7
All Michigan Counties	3,982,000	3.4	1367

*Source: MESC 1993 Annual Average Labor Statistics for Employment by Place of Work. Some employee population data is only available at a multi-county level, as indicated (i.e. not available at a single county level). Therefore, some data is presented with grouped counties. **Rates are based on the average number of cases per year from 1989-1998, per 100,000 Michigan workers. ***MSA=Metropolitan Statistical Area and includes Lapeer (16 cases), Macomb (138 cases), Monroe (9 cases), Oakland (189 cases), St. Clair (10 cases) and Wayne (376 cases) counties.

Table 3. Annual Incidence Rates of Work-Related Asthma Among Michigan Workers by Major Metropolitan Area: 1990-1998

	<u>Clinton-Eaton-Ingham</u>	<u>Kent-Ottawa-Muskegon-Allegan</u>	<u>Saginaw-Bay-Midland</u>	<u>Detroit MSA**</u>	<u>Total (all Michigan)</u>
1990	1.4 (3)	2.2 (8)	2.4 (4)	3.0 (58)	3.6 (144)
1991	3.8 (8)	1.4 (5)	4.3 (7)	2.7 (50)	2.9 (114)
1992	5.6 (12)	0.7 (3)	1.8 (3)	4.6 (86)	3.8 (150)
1993	3.7 (8)	1.3 (6)	1.8 (4)	6.4 (121)	4.4 (176)
1994	1.8 (4)	3.5 (7)	1.8 (3)	4.4 (85)	3.7 (152)
1995	2.2 (5)	1.2 (6)	1.7 (3)	3.4 (69)	2.9 (122)
1996	1.3 (3)	0.9 (5)	2.9 (5)	4.0 (91)	3.4 (149)
1997	2.2 (5)	1.1 (6)	4.5 (8)	3.7 (77)	3.5 (156)
1998	2.6 (6)	1.4 (8)	3.9 (7)	3.7 (79)	3.1 (141)

*Rate per 100,000 Michigan workers. Rate, number of cases in parentheses.

Source: MESC Annual Average Labor Statistics for Employment by Place of Work, for each year 1990-1998 separately.

**MSA=Metropolitan Statistical Area. For the years 1990-1995, includes Lapeer, Livingston, Macomb, Monroe, Oakland, St. Clair, and Wayne counties. For 1996, 1997, and 1998, does not include Livingston county because of a change in the counties associated with certain MSA's (including Detroit), beginning in 1996.

**Table 4. Primary Industrial Exposure for Confirmed Work-Related
Asthma Patients: 1988-2000**

<u>Industry (SIC Code)*</u>	<u>Number of Cases</u> <u>1988-2000**</u>		<u>Number of</u> <u>Employees***</u>	<u>Ann. Average</u> <u>Inc.</u> <u>Rate****</u> <u>1989-1998</u>	
MANUFACTURING (20-39)					
Automobile (37)	710	(43.3)	279,000	20.9	(583)
Ind. & Comm. Mach. & Computer Equipment (35)	63	(3.8)	117,000	4.9	(57)
Fabricated Metal Products (34)	72	(4.4)	116,000	5.8	(67)
Rubber and Misc. Plastic Products (30)	65	(4.0)	55,000	10.5	(58)
Foundries (33)	55	(3.4)	36,000	13.3	(48)
Food and Kindred Products (20)	31	(1.9)	44,000	6.6	(29)
Printing and Publishing (27)	16	(1.0)	45,000	2.9	(13)
Paper and Allied Products (26)	15	(0.9)	21,000	6.2	(13)
Lumber and Wood (24)	16	(1.0)	15,000	8.0	(12)
Electrical Equipment (36)	13	(0.8)	30,000	4.0	(12)
Furniture and Fixtures (25)	8	(0.5)	34,000	1.5	(5)
Apparel Made from Fabric (23)	2	(0.1)	18,000	1.1	(2)
Other Durables (32,38,39)	38	(2.3)	41,000	7.8	(32)
Other Nondurables (22,28,29,31)	67	(4.1)	50,000	11.4	(57)
WHOLESALE AND RETAIL TRADE (50-59)					
Wholesale-Nondurable Goods (51)	11	(0.7)	72,000	1.4	(10)
Food Stores (54)	11	(0.7)	101,000	1.1	(11)
Eating and Drinking Places (58)	8	(0.5)	262,000	0.3	(7)
Wholesale-Durable Goods (50)	11	(0.7)	125,000	0.9	(11)
Automotive Dealers and Gasoline Services (55)	8	(0.5)	76,000	0.5	(4)
General Merchandise Stores (53)	4	(0.2)	120,000	0.3	(4)
Miscellaneous Retail (52, 56, 57,59)	6	(0.4)	141,000	0.3	(4)
SERVICES					
Health (80)	147	(9.0)	331,000	3.6	(119)
Education (82)	51	(3.1)	38,000	11.8	(45)
Automotive Repair (75)	10	(0.6)	34,000	2.4	(8)
Business (73)	13	(0.8)	203,000	0.6	(12)
Social Services (83)	7	(0.4)	74,000	0.9	(7)
Engineering, Accounting, etc. (87)	6	(0.4)	83,000	0.7	(6)
Other Services (70,72,76,79,81,86,89)	38	(2.3)	254,000	1.3	(34)
CONSTRUCTION AND MINING (10-17)					
Special Trade Construction (17)	30	(1.8)	89,000	2.9	(26)
Other Construction (15-16)	8	(0.5)	43,000	1.9	(8)
Mining (10-14)	4	(0.2)	9,000	4.4	(4)
MISCELLANEOUS INDUSTRIES					
Government (91-97)	51	(3.1)	640,000	0.7	(46)
Transportation and Utilities (40-49)	26	(1.6)	156,000	1.5	(23)
Finance, Insurance and Real Estate (60-67)	8	(0.5)	192,000	0.4	(8)
Agricultural Production and Services (01,02,07)	5	(0.3)	90,000	0.3	(3)
Unknown	4	(0.2)	—	—	(3)
TOTAL	1638	(100.0)	3,982,000	3.5	(1388)

*Standard Industrial Classification.

**Number of cases, percentages are in parentheses.

***Source:MESC 1993 civilian labor force and industrial employment estimates.

****Average annual incidence rate, total number of cases for 1989-1998 are in parentheses. Rates are based on average number of cases from 1989-1998 per 100,000 adult workers in each industrial category.

**Table 5. Primary Industrial Exposure for Confirmed
Work-Related Asthma Patients: 1990-1998**

<u>INDUSTRY (SIC code*)</u>	<u>1990**</u>	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>
Manufacturing (20-39)									
Food & Kindred Products (20)	20.0 (9)	6.7 (3)	2.3 (1)	-- (0)	2.3 (1)	4.4 (2)	6.8 (3)	4.9 (2)	7.5 (3)
Chemicals & Allied Product & Petroleum Refining (28,29)	7.0 (8)	4.3 (2)	14.0 (7)	12.0 (6)	7.8 (4)	14.3 (7)	*** (3)	*** (7)	*** (5)
Rubber & Misc. Plastic Products (30)	9.3 (5)	19.6 (10)	-- (0)	7.3 (4)	11.3 (7)	15.1 (10)	10.4 (7)	13.2 (9)	4.5 (3)
Foundries (33)	9.5 (4)	23.7 (9)	8.1 (3)	19.4 (7)	18.9 (7)	5.4 (2)	18.9 (7)	8.1 (3)	10.5 (4)
Fabricated Metal Products (34)	6.6 (8)	4.4 (5)	2.6 (3)	3.4 (4)	5.7 (7)	3.1 (4)	12.5 (16)	8.6 (11)	4.7 (6)
Industrial & Commercial Machinery & Computer Equipment (35)	6.3 (8)	1.7 (2)	6.1 (7)	8.5 (10)	6.6 (8)	3.0 (4)	3.0 (4)	2.2 (3)	3.7 (5)
Automobile (37)	19.0 (56)	15.6 (44)	28.7 (82)	26.5 (74)	25.0 (71)	17.3 (51)	21.1 (59)	21.1 (61)	23.7 (65)
Other Durables (38,39)	19.2 (5)	8.0 (2)	4.9 (2)	9.8 (4)	-- (0)	-- (0)	6.7 (3)	4.4 (2)	6.5 (3)
Miscellaneous Industries									
Special Trade Construction (17)	1.1 (1)	2.3 (2)	4.6 (4)	6.7 (6)	3.2 (3)	0.9 (1)	1.7 (2)	0.8 (1)	2.4 (3)
Transportation & Utilities (40-49)	0.6 (1)	1.3 (2)	1.9 (3)	4.5 (7)	2.5 (4)	1.2 (2)	B (0)	0.6 (1)	0.6 (1)
Health (80)	1.6 (5)	1.9 (6)	3.4 (11)	3.6 (12)	3.8 (13)	3.7 (13)	3.3 (12)	6.3 (23)	6.0 (22)
Education (82)	5.4 (2)	5.3 (2)	8.1 (3)	21.1 (8)	7.7 (3)	10.2 (4)	18.6 (8)	20.5 (9)	11.4 (5)
Government (91-97)	8.1 (5)	6.6 (4)	7.9 (5)	14.3 (9)	6.3 (4)	1.1 (7)	0.8 (5)	0.5 (3)	0.5 (3)
Total (all industries)	3.4 (144)	2.8 (114)	3.5 (150)	4.0 (176)	3.4 (152)	3.0 (127)	3.3 (152)	3.2 (161)	3.0 (147)

*Standard Industrial Classification.

**Annual Incidence rate; number of cases for that year are in parentheses. Rates are based on the number of cases per 100,000 adult workers in Michigan for each year separately.

Source: MESC Annual Average civilian labor force and industrial employment estimates.

***Denominator not available in 1996, 1997 and 1998.

**Table 6. Occupational Agents Associated with 1638 Confirmed
Work-Related Asthma Patients: 1988-2000**

<u>Allergen</u>	<u>Number</u>	<u>Percent</u>
Isocyanates	295	18.0
Metal-Working Fluids	198	12.1
Unknown (Mfg.)	107	6.5
Unknown (Office)	100	6.1
Cleaning Solutions	93	5.7
Exhaust, Smoke, Fumes	88	5.4
Welding Fumes	82	5.0
Solvent	49	3.0
Latex/Rubber	49	3.0
Formaldehyde	40	2.4
Epoxy	40	2.4
Paint Fumes	36	2.2
Acrylates	30	1.8
Acids	30	1.8
Chlorine	26	1.6
Cobalt	21	1.3
Styrene	20	1.2
Wood Dust	19	1.2
Plastic Fumes	15	0.9
Flour	13	0.8
Ammonia	12	0.7
Chromium	11	0.7
Herbicide, Pesticide	11	0.7
Printing Inks	11	0.7
Amines	9	0.5
Caustics	8	0.5
Animals	8	0.5
Grain Dust	7	0.4
Fiberglass	7	0.4
Meat Wrappers' Asthma	7	0.4
Pickling Ingredients	6	0.4
Cement Dust	5	0.3
Rose Hips	4	0.2
Sulfonate	4	0.2
Sulfur Dioxide	4	0.2
1,1,1 Trichloroethane	4	0.2
Cosmetology Chemicals	4	0.2
Nitrogen	4	0.2
Glutaraldehyde	4	0.2
Colophony	3	0.2
Photo Developing Fluids	3	0.2
Phthalic Anhydride	3	0.2
Cadmium Solder	2	0.1
Enzymes	2	0.1
Nickel	2	0.1
Other*	142	8.7
Total	1638	99.6**

*Includes: x-ray developing fluids, chemicals used in the construction industry, freon, sulfite, gas and oil refinery exposures, maleic anhydride, solder fumes, vinyl acetate, ozone, zinc, textile lint, 1,3, dichloro-2-propanol, pepper gas, exercise, weeds, soda ash, asbestos, limone, coal dust, ethylene oxide, trichloroethylene, iodine, polyvinyl butyrate, lime dust, polyester, MEK, nylon-polyhexamethylene adipamide, potassium aluminum fluoride, ammonium hydroxide, methanol, naphtha, drywall dust, paper dust, zinc oxide, methylene chloride, ethylene glycol monobutyl ether, sewage, tar fumes, monoammonium phosphate, potassium hydroxide, ethyl alcohol, heptane, stress, sodium hydroxide, platinum, phosgene, cyanide, heat, polyethylene, perchloroethylene, hydraulic oil, blood, 1,3 dichloro 5 5-dimethyl hydrantoin, ninhydrin, cigarette smoke, azodicarbamide, lactase, medications, sodium acetate, cellulose, glaze, sulfuric acid, aspirin, tetrahydrofuran, zinc borate, leather, rust inhibitor, citrus spray, mold release, sand, hydrogen sulfide, teflon, flux, natural gas, tuberculosis vaccine, fungus, gortex, hydrogen peroxide, sludge, and premicide.

**Percentages do not add to 100 due to rounding.

**Table 7. Cigarette Smoking Status of Confirmed
Work-Related Asthma Patients: 1988-2000**

<u>Smoking Status</u>	<u>Disease Status</u>				
	<u>ALL*</u>	<u>OA**</u>	<u>POA</u>	<u>AA</u>	<u>RA</u>
Current Smoker	318 (19.9)	160 (22.5)	93 (16.0)	33 (19.8)	32 (22.5)
Ex-Smoker	659 (41.2)	280 (39.4)	256 (44.1)	59 (35.3)	64 (45.1)
Non-Smoker	623 (38.9)	270 (38.0)	232 (39.9)	75 (44.9)	46 (32.4)
Total	1600	710	581	167	142

*Total number of cases: 1600. Smoking status was missing on 38 individuals. Number of patients, percentages are in parentheses.

**OA=occupational asthma; POA=possible occupational asthma; AA=aggravated asthma; RA=reactive airway dysfunction syndrome.

Table 8. Family History of Allergies Among Confirmed Work-Related Asthma Patients: 1988-2000

Family History of Allergies	Disease Status				
	<u>ALL</u>*	<u>OA</u>**	<u>POA</u>	<u>AA</u>	<u>RA</u>
YES	615 (42.0)	257 (39.0)	221 (41.3)	85 (59.9)	52 (40.9)
NO	848 (58.0)	402 (61.0)	314 (58.7)	57 (40.1)	75 (59.1)
Total	1463	659	535	142	127

*Total number of cases: 1463. Missing data on 175 patients. Number of patients, percentages are in parentheses.

**OA=occupational asthma; POA=possible occupational asthma; AA=aggravated asthma; RA=reactive airway dysfunction syndrome.

**Table 9. Personal History of Allergies or Asthma
Among Confirmed Work-Related Asthma
Patients: 1988-2000**

<u>Personal History</u>	Disease Status				
	<u>ALL*</u>	<u>OA**</u>	<u>POA</u>	<u>AA</u>	<u>RA</u>
YES	737 (45.0)	284 (39.3)	245 (41.2)	159 (89.8)	49 (33.8)
NO	901 (55.0)	438 (60.7)	349 (58.8)	18 (10.2)	96 (66.2)
Total	1638	722	594	177	145

*Number of patients, percentages are in parentheses.

**OA=occupational asthma; POA=possible occupational asthma; AA=aggravated asthma; RA=reactive airway dysfunction syndrome.

**Table 10. Persistence of Symptoms and Medication
Use in Confirmed Work-Related Asthma Patients: 1988-2000**

<u>Exposure Status</u>	<u>Total*</u>	<u>Symptoms</u>		<u>Asthma Medications</u>	
		<u>Yes</u>	<u>Less</u>	<u>Yes</u>	<u>Less</u>
Still Exposed	464	442 (95.3)	154 (33.2)	392 (84.5)	98 (21.1)
No Longer Exposed	1057	884 (83.6)	528 (50.0)	810 (76.6)	316 (29.9)
<hr/>					
Total	1521	1326	682	1202	414

*Total number of cases: 1521. Information missing on 117 individuals. Number of patients, percentages are in parentheses.

**Table 11. Status of Facilities Where 1638 Patients with
Confirmed Work-Related Asthma were
Exposed to Allergens: 1988-2000**

<u>Inspection Status</u>	<u>Number of Patients Represented</u>	<u>Number</u>	<u>Percent</u>
Inspections	881	513*	44.6
No Follow-up Planned	630	519	45.1
Scheduled for Inspection	3	3	0.3
Closed	35	33	2.9
No Longer Use Occupational Allergen	22	21**	1.8
Sent Company an Indoor Air Letter	20	16	1.4
Sent Company a Letter to Check Exposures	47	46	4.0
Total	1638	1151***	100.1****

*513 inspections were conducted in 451 different facilities.

**Five companies that no longer use the allergen were previously inspected.

***Represents 1,089 different facilities.

****Percent does not add to 100 due to rounding.

**Table 12. Results of 513 Industrial Hygiene Inspections in
451 Facilities Where Patients with Confirmed
Work-Related Asthma were Exposed to Allergens: 1988-2000**

<u>Inspection Results</u>	<u>Number</u>	<u>Percent</u>
Air Sampling - NIOSH Standard		
Above NIOSH Standard	49	9.6
Below NIOSH Standard	304	59.3
No NIOSH Standard	9	1.8
Unknown (no report yet)	5	1.0
Did Not Sample for an Allergen	9	1.8
<u>Did Not Sample</u>	<u>137</u>	<u>26.7</u>
Total	513	100.2*
Air Sampling - MIOSHA Standard		
Above MIOSHA Standard	24	4.7
Below MIOSHA Standard	337	65.7
No MIOSHA Standard	2	0.4
Unknown (no report yet)	5	1.0
Did Not Sample for an Allergen	8	1.6
<u>Did Not Sample</u>	<u>137</u>	<u>26.7</u>
Total	513	100.1*

*Percent does not add to 100 due to rounding.

**Table 13. Symptoms Consistent with Work-Related Asthma
Among Fellow Workers of the 1638 Confirmed
Work-Related Asthma Patients**

<u>Symptoms*</u>	Disease Status of Index Patient				
	<u>ALL**</u>	<u>OA</u>	<u>POA</u>	<u>AA</u>	<u>RA</u>
Daily or Weekly SOB, Wheezing or Chest Tightness	1257 (17.1)	943 (17.8)	279 (16.0)	4 (7.5)	31 (13.5)
<u>OSHA Log***</u>	430 (17.5)	314 (21.1)	105 (11.5)	2 (12.5)	9 (11.1)
Total	1687****	1257***	384	6	40

*Denominator for calculating percentages was the number of workers interviewed. SOB=shortness of breath.

**Number of individuals with symptoms, percentages are in parentheses. OA=occupational asthma; POA=possible occupational asthma;
AA=aggravated asthma; RA=reactive airways dysfunction syndrome.

***Numerator for calculating percentages was the number of companies with an employee other than the index patient on the OSHA log.
Denominator for calculating percentages was the number of companies inspected.

****Eight individuals were identified both on the questionnaire and the OSHA log.